## Open-Channel Hydraulics H.W.#3 Reynolds Equations

Professor Sung-Uk Choi Mar.26, 2025

- 1. Express  $\nabla V^2/2$  and  $\overline{V \times curl V}$  in terms of u, u', v, v', w, and w' for turbulent flows.
- 2. Express the average ratio of dilatational and shear deformation in terms of mean and fluctuating velocity components for a turbulent flows.
- 3. Demonstrate that the Reynolds equations can still be written:

$$\frac{\partial \overline{u}}{\partial t} + \overline{u} \frac{\partial \overline{u}}{\partial x} + \overline{v} \frac{\partial \overline{u}}{\partial y} + \overline{w} \frac{\partial \overline{u}}{\partial z} = \frac{\partial}{\partial x} \left[ \frac{p^*}{\rho} + 2v \frac{\partial \overline{u}}{\partial x} - \overline{u'^2} \right] 
+ \frac{\partial}{\partial y} \left[ v \left( \frac{\partial \overline{u}}{\partial y} + \frac{\partial \overline{v}}{\partial x} \right) - \overline{u'v'} \right] 
+ \frac{\partial}{\partial z} \left[ v \left( \frac{\partial \overline{u}}{\partial z} + \frac{\partial \overline{w}}{\partial x} \right) - \overline{u'w'} \right]$$

and two other equations which will be determined. Indicate the advantage of this form of the Reynolds equations.

4. Write the Reynolds equations in the case of a mean two-dimensional motion. Write the Reynolds equations in the case of isotropic turbulence, i.e.,  $u'^2 = v'^2$  and u'v' = 0.